## **Immunizations, adults 65+**

Health Priority E: Existing, Emerging, and Re-emerging Communicable Diseases

Objective E2c: By 2010, at least 90 percent of Wisconsin residents 65 years of age and older and individuals with chronic health conditions will be fully immunized in accordance with current Advisory Committee on Immunization Practices (ACIP) recommendations.

2010 Target: Greater than or equal to 90%

Indicator: Immunizations, adults 65+

## Percent of Wisconsin Adults Aged 65 and Older Who Are Immunized in Accordance with ACIP Recommendations

initialized in Accordance with ACII Recommendations									
	Flu Shot	t in Past	Pneumor	nia Shot					
Year	12 Mc	onths	(Ev	er)	Both				
(N)	Percent	+/-	Percent	+/-	Percent	+/-			
2001	70%	4%	65%	4%	55%	4%			
(N)	590		590		590				
2002	74%	3%	71%	3%	61%	3%			
(N)	847		829		829				
2003	72%	3%	67%	3%	58%	3%			
(N)	805		783		783				
2004	74%	3%	70%	4%	62%	4%			
(N)	921		895		895				
2005	72%	3%	66%	4%	56%	4%			
(N)	1,121		1,083		1,082				

Source: Wisconsin Behavioral Risk Factor Survey, Bureau of Health Information and Policy, Division of Public Health, Department of Health and Family Services. Note: ACIP also recommends tetanus and diphtheria boosters for adults aged 65+. Data on these immunizations are not reported in available sources.

- N Number in sample. See data documentation.
- +/- Confidence interval. See data documentation

**Continued on next page** 

Prepared for Tracking the State Health Plan, 2010—State-Level Data <a href="http://dhfs.wisconsin.gov/statehealthplan/track2010">http://dhfs.wisconsin.gov/statehealthplan/track2010</a> (July 2006)
Bureau of Health Information and Policy
Division of Public Health
Wisconsin Department of Health and Family Services

## Immunizations, adults 65+, continued

## Percent of Wisconsin Adults Aged 65 and Older Who Are Immunized in Accordance with ACIP Recommendations, by Sex

	Flu Shot in				Pneumonia Shot							
	P	months	(Ever)			Both						
Year	Males		Females		Males		Females		Males		Females	
( <b>N</b> )	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-
2001	75%	6%	67%	5%	63%	7%	67%	5%	56%	7%	56%	5%
(N)	204		384		196		379		196		379	
2002	76%	5%	73%	4%	68%	6%	72%	4%	60%	6%	61%	4%
(N)	281		566		267		562		267		562	
2003	71%	5%	73%	4%	62%	6%	70%	4%	54%	6%	60%	4%
(N)	285		520		272		511		272		511	
2004	76%	5%	73%	4%	67%	6%	73%	5%	61%	6%	62%	5%
(N)	310		611		298		597		298		597	
2005	74%	5%	71%	5%	64%	6%	67%	4%	57%	6%	55%	5%
(N)	398		723		376		707		376		706	

Source: Wisconsin Behavioral Risk Factor Survey, Bureau of Health Information and Policy, Division of

Public Health, Department of Health and Family Services.

Note: ACIP also recommends tetanus and diphtheria boosters for adults aged 65+. Data on these immunizations are not reported in available sources.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation